

Spontaneous Nephrocutaneous fistula due to Xanthogranulomatous pyelonephritis with secondary Enterocutaneous fistula: A rare case report

Reza kazemi

University of Medical Sciences, Isfahan, Iran.

Abstract

Chronic infection, especially in the setting of obstruction like calculous disease, may result in the fistula tract development of fistula tract from kidney to the other proximal organs. However, nephrocutaneous fistula is a rare complication, and the primary cause of its development is previous surgical intervention, trauma, or malignancy. Our case was a 26-year-old Afghan man with xanthogranulomatous pyelonephritis (XPN) resulted in spontaneous nephrocutaneous fistula. The patient underwent radical nephrectomy. Our case shows that previous surgery is not the only cause of fistula. After surgery, he presented with secondary enterocutaneous fistula due to surgery or chronic underlying inflammation. After systemic antibiotic therapy and total parenteral nutrition, he became well and discharged from the hospital. As a result, XPN is a rare subtype of chronic pyelonephritis that requires immediate evaluation and early diagnosis. Patients should be considered for possible complications such as fistulas. Physicians should be aware of this issue for appropriate diagnosis and treatment.

Received: February 05, 2022; **Accepted:** February 08, 2022; **Published:** February 11, 2022

Biography

Reza kazemi has completed his General Medicine at the age of 25 years from Islamic Azad University, Najafabad Branch, and Urology specialty studies from Isfahan University of Medical Sciences and He also holds the top rank in the Urology Board certified. He is

Assistant Professor, Department of Urology, Al-Zahra Hospital, Isfahan University of Medical Sciences, Isfahan, Iran. He has published more than 9 papers in journals and He is currently working on several research projects.